

Lecture 39  
23 April 2008

Vertebrate Physiology  
ECOL 437 (MCB/VetSci 437)  
Univ. of Arizona, spring 2008

Kevin Bonine & Kevin Oh



## 1. Metabolism (Ch5)

[http://eebweb.arizona.edu/eeb\\_course\\_websites.htm](http://eebweb.arizona.edu/eeb_course_websites.htm)

1

Housekeeping, 23 April 2008



---

### Upcoming Readings

Wed 23 Apr: Ch 4, 5

LAB 23 Apr: Kevin Oh emailed

Final Proposal due in lab 23 April or  
beginning of lecture 25 April

Fri 25 Apr: Ch 4,5

---

Lab discussion leaders: 23 April

1pm - none

3pm - Nina


2

## Metabolism



ANIMAL PHYSIOLOGY, Figure 7.2 © 2004 Sinauer Associates, Inc.

## World Class Human Runners



ANIMAL PHYSIOLOGY, Figure 6.9 © 2004 Sinauer Associates, Inc.

(Hill et al. 6.9)

4

## Metabolism

- Chemical reactions in the body
- Temperature**-dependent rates
- Not 100% efficient, **energy lost as heat**  
(not 'lost' if used to maintain Tb)

### 1. Anabolic


- creation, assembly, repair, **growth**  
(positive nitrogen balance)

### 2. Catabolic

- energy release** from complex molecules  
(carbos, fats, proteins)
- energy storage** in **phosphate bonds** (ATP) and  
**metabolic intermediates** (glucose, lactate)


5

## Chemical Energy



6

## 10% Rule



Hill et al. 2004, Fig 4.9

ANIMAL PHYSIOLOGY, Figure 4.9 © 2004 Sinauer Associates, Inc.

## Metabolism


### Energy Available for:

- Growth, Maintenance, Reproduction

- SDA (specific dynamic action)


*No free lunch!*

(b) SDA in flatfish



ANIMAL PHYSIOLOGY, Figure 5.5 (Part B) © 2004 Sinauer Associates, Inc.

(a) The concept of SDA




(Hill et al. 5.5)

ANIMAL PHYSIOLOGY, Figure 5.5 (Part A) © 2004 Sinauer Associates, Inc.

## Metabolism and Ecology

Male emperor penguin  
>100days w/o food when  
incubating eggs



Knut Schmidt\_Nielsen 1997 9

## Metabolism and Ecology

K vs. r selected  
(logistic curve)


-Larger animals invest  
proportionally less in  
reproduction

-Sperm is cheap

-Direct and indirect costs



## Fetal teeth in caecilians



## Metabolic Rate

-measurable conversion of chemical energy into heat

-used to understand:

- energy budgets
- dietary needs
- body size implications
- habitat effects
- costs of various activities
- mode of locomotion
- cost of reproduction

Much more difficult for **water breathing animals** to maintain body temperatures above ambient because **rate of heat transfer is greater than rate of O<sub>2</sub> transfer** in water (high specific heat)



*Amblyrhynchus cristatus*



**SEAL BLUBBER** This cross section of a frozen seal shows the thick layer of blubber. Of the total area in the photo, 58% is blubber and the remaining 42% is muscle, bone, and visceral organs. The measuring stick is graduated in inches. [Courtesy of P. F. Scholander, University of California, San Diego]  
Knut Schmidt\_Nielsen 1997


13

Fish Example:


-Differences in **vascular organization**

-Tuna with **warm, aerobic muscle** medially

-**Countercurrent** blood flow (don't lose heat to cold water across gills)







**HEAT EXCHANGER** Cross section of a 2 kg skip-jack tuna (*Katsuwonis pelamis*) shows how this powerful swimming machine consists mostly of muscle (left). The red muscle, which is maintained at high temperature, appears nearly black in the photo. In some tunas the heat exchanger is located laterally, but in the skipjack the main heat exchanger is located just below the vertebral column, almost exactly in dead center of the photo. A cross section

of the vascular heat exchanger (right) shows a roughly equal number of arteries and veins. The arteries (smaller and thick-walled) are interspersed with veins (larger and thin-walled). The diameter of the arteries is about 0.04 mm and of the veins 0.08 mm, and their length is about 10 mm. [Courtesy of E. D. Stevens, University of Guelph, Ontario]

Knut Schmidt\_Nielsen 1997

## Metabolic Rates

### -Basal Metabolic Rate, BMR

- minimal environmental and physiological stress (appropriate ambient temperature, post-digestive, resting etc.)

### -Standard Metabolic Rate, SMR

- similar to BMR, but at a given  $T_b$

### -Field Metabolic Rate, FMR

- average metabolic rate of animal in natural setting
- hard to measure







Figure 7.11 Body temperature of a willow tit (*Parus montanus*) during the night hours at three different ambient temperatures. The records are from midafternoon one day to the following morning. [Reinertsen and Haftorn 1986]

Knut Schmidt\_Nielsen 1997



Knut Schmidt\_Nielsen 1997

17

## Metabolic Rates

### Basal Metabolic Rate, BMR


-important components:

1. Membrane form and function  
 maintenance of electrochemical gradients  
 -proton pumps in mitochondrial membranes  
 -Na/K-ATPase pumps in plasma membrane
2. Protein synthesis
3. ATP formation


18

Metabolic heat production  
(chemical energy 'lost' as heat during metabolism)

- Endotherms
- surface area to volume ratio
- Larger ectotherms can be heterothermic
  - leatherback (*Dermochelys coriacea*)
  - pythons (female brooding clutch)
  - tuna and increased core temperature



(a)




(b)

Pough et al., 2001


## Digestive Systems

Transit time (time to digest),  
cost, and anatomy variable:


- Food quality
- Body Size
- Temperature (ectotherms)




(a)



(Eckert 15-12)



(Eckert 15-11)



(Eckert 15-16)

Gut Plasticity

Alter gut [size](#), [activity](#) ([reversible](#))

-Sustained increased metabolism can increase bird [gut length](#) by 1/5



© 2002 Michael Porter  
Click to enlarge


-Mammals increase [GI tract mass](#) 3-4x post-hibernation



© BMC 2000

## Gut Plasticity

Alter gut size, activity (reversible)



-Some infrequently-feeding snakes:

- intestine 2x larger within 2 days
- microvilli length and area up 400%
- glucose transport rate up as much as 22x
- other transporters also up-regulated (e.g., a.a. absorption)



Alkaline Tide...